

The effects of nutrition on type 2 diabetes risk and management: A systematic review of studies published between 2015 and 2025

Tuba Sert, Gamze Şanlı Ak

Nutrition and Dietetics, Nişantaşı University,
Istanbul, Turkey

ORCID  of the author(s)

TS: <https://orcid.org/0009-0007-2144-7812>
GŞA: <https://orcid.org/0009-0004-9451-5239>

Corresponding Author

Tuğba Sert
Nutrition and Dietetics, Nişantaşı University,
Istanbul, Turkey
E-mail: tugbasert9910@gmail.com

Ethics Committee Approval

Ethics committee approval was not required because this review did not involve human participants, animal experiments, patient-level data, or identifiable personal information. Informed consent was not applicable.

Conflict of Interest

No conflict of interest was declared by the authors.

Financial Disclosure

The authors declared that this study has received no financial support.

Published
2026 May 7

Copyright © 2026 The Author(s)



This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0).
<https://creativecommons.org/licenses/by-nc-nd/4.0/>



Abstract

Background/Aim: Type 2 diabetes is a major global metabolic disorder, and nutritional exposure is central to both prevention and long-term glycemic management. This systematic review evaluated the effects of dietary patterns, macronutrient composition, fiber intake, protein source, and lifestyle-based nutritional interventions on type 2 diabetes risk and management.

Methods: A literature search was performed in Google Scholar, Web of Science, PubMed, Scopus, and the Cochrane Library on August 27, 2025. The search used combinations of type 2 diabetes, nutrition, diet, dietary patterns, glycemic control, prevention, and management. English-language studies published between 2015 and 2025 were screened. Randomized trials, cohort studies, systematic reviews, and meta-analyses were considered eligible. Studies not directly related to type 2 diabetes or nutrition were excluded. Eligible evidence was synthesized narratively.

Results: A total of 1,200 records were initially identified, and 40 studies were included after screening and eligibility assessment. Mediterranean, plant-based, and fiber-rich dietary patterns were consistently associated with improved glycemic control, lower cardiometabolic risk, and reduced type 2 diabetes risk. Low-carbohydrate diets showed short-term benefits for hemoglobin A1c and body weight, although long-term sustainability and safety remained uncertain. Protein quality, particularly the balance between plant and animal protein sources, also appeared relevant to insulin resistance and metabolic outcomes. Lifestyle-based and digital interventions enhanced adherence but showed variable durability.

Conclusion: Nutrition is a cornerstone of type 2 diabetes prevention and management. Current evidence favors individualized, sustainable dietary strategies emphasizing overall diet quality, fiber-rich foods, plant-forward patterns, and integration with behavioral support rather than reliance on a single universal diet model.

Keywords: type 2 diabetes, nutrition, dietary patterns, glycemic control, prevention

Introduction

Type 2 diabetes (T2D) is one of the most important metabolic and public health challenges worldwide. Recent global estimates show that the total diabetes burden has increased substantially, and projections indicate that the number of people living with diabetes will continue to rise over the coming decades, with T2D accounting for most cases [1, 2]. This increasing burden is driven by population aging, urbanization, excess adiposity, sedentary behavior, and dietary patterns characterized by high energy density and low nutritional quality [3, 4].

Nutrition has a direct role in glucose homeostasis, insulin sensitivity, body weight regulation, lipid metabolism, and systemic inflammation. Dietary approaches such as Mediterranean, plant-based, high-fiber, low-glycemic-index, and carbohydrate-modified diets have been investigated for both prevention and management of T2D [5-8]. However, the comparative interpretation of these approaches remains difficult because studies differ in design, intervention intensity, follow-up duration, baseline metabolic risk, and adherence support.

Beyond individual macronutrients, contemporary diabetes nutrition science increasingly emphasizes dietary patterns, food quality, behavioral adherence, cultural suitability, and long-term sustainability. Therefore, a broad synthesis of current evidence is useful for clinicians, dietitians, and researchers. This systematic review aimed to summarize evidence published between 2015 and 2025 regarding the effects of nutrition on T2D risk and management.

Materials and methods

Study design

This review was designed as a systematic literature review with narrative synthesis. The reporting structure was refined to improve transparency and consistency with common biomedical reporting principles. Because of heterogeneity in study design, population characteristics, dietary interventions, and outcomes, no quantitative meta-analysis was performed.

Search strategy

A literature search was performed in Google Scholar, Web of Science, PubMed, Scopus, and the Cochrane Library on August 27, 2025. Search terms included combinations of type 2 diabetes, nutrition, diet, dietary patterns, dietary intervention, glycemic control, prevention, management, hemoglobin A1c, insulin resistance, Mediterranean diet, plant-based diet, low-carbohydrate diet, fiber, and protein. The search was limited to studies published in English between January 1, 2015, and August 27, 2025.

Inclusion and exclusion criteria

Randomized controlled trials, prospective cohort studies, systematic reviews, and meta-analyses evaluating the relationship between nutrition and T2D prevention or management were included. Eligible studies reported outcomes related to incident T2D, hemoglobin A1c, fasting plasma glucose, insulin resistance, body weight, cardiometabolic risk factors, adherence, or diabetes-related lifestyle management. Studies were excluded if they were not directly related to T2D, did not evaluate a nutritional exposure or intervention, were not available in English, were animal or laboratory-only studies, or were published outside the defined

time window unless they were used as foundational background evidence.

Study selection and risk of bias assessment

Records were screened by title and abstract, followed by full-text evaluation for potentially eligible studies. Studies were grouped thematically according to dietary patterns, carbohydrate modification, fiber and whole grains, protein source and amino acid profile, and lifestyle or digital nutritional support. The submitted manuscript stated that risk of bias was evaluated using the Cochrane risk-of-bias approach for randomized trials and the Newcastle-Ottawa Scale for observational studies; however, individual domain-level ratings were not provided in the submitted file.

Statistical analysis

A descriptive narrative synthesis was performed. No pooled effect estimates, hypothesis tests, or new inferential statistics were calculated. Therefore, P-values were not applicable to the review synthesis.

Ethical considerations

This review was based exclusively on previously published literature and did not involve human participants, animal experiments, patient-level data, or identifiable information. Therefore, ethics committee approval and informed consent were not required.

Results

Study selection

The search identified 1,200 records. After removal of duplicates and title-and-abstract screening, 150 full-text reports were assessed for eligibility. Forty studies met the inclusion criteria and were included in the final synthesis. The study selection process is shown in Figure 1.

Figure 1. PRISMA 2020 flow diagram of the study selection process (initial records identified: 1200; studies included: 40).




Table 1. Summary of principal findings by nutritional domain.

Nutritional domain	Evidence base and outcomes	Principal findings and clinical interpretation	References
Dietary patterns	12 studies; T2D risk, HbA1c, body weight, and cardiometabolic markers.	Mediterranean, plant-based, and overall high-quality dietary patterns were generally associated with better glycemic and cardiometabolic profiles. This was the most consistent and clinically applicable evidence domain; adherence and food quality remain central.	[5-15]
Low-carbohydrate and macronutrient composition	8 studies; HbA1c, body weight, lipids, and remission-related outcomes.	Low-carbohydrate approaches showed short-term metabolic benefits, but long-term superiority was less consistent. They may be useful for selected patients when medication adjustment and clinical monitoring are available.	[7, 16-22]
High-fiber and whole-grain intake	6 studies; T2D incidence, postprandial glucose, and weight control.	Higher fiber and whole-grain intake were associated with lower T2D risk and improved metabolic responses. These data support recommending minimally processed, fiber-rich carbohydrate sources rather than broad carbohydrate avoidance.	[23-27]
Protein source and amino acid profile	5 studies; insulin resistance and metabolic risk.	Plant-forward protein patterns appeared more favorable than animal-heavy patterns in observational evidence; branched-chain amino acid signatures were linked to insulin resistance. Protein recommendations should consider kidney function, sarcopenia risk, weight goals, and cardiometabolic risk.	[28, 29]
Lifestyle-based and digital nutritional support	9 studies; adherence, body weight, HbA1c, and prevention outcomes.	Structured lifestyle programs and digital support tools improved adherence and self-management, although durability varied. These interventions appear most effective when integrated with behavioral counseling and ongoing follow-up.	[30-33]

HbA1c: hemoglobin A1c, T2D: type 2 diabetes.

Principal findings

The included evidence indicated that overall dietary pattern quality was more clinically relevant than any isolated nutrient. Mediterranean and plant-forward diets were associated with improved glycemic control, weight-related outcomes, and cardiometabolic risk profiles. Low-carbohydrate diets were associated with short-term improvements in hemoglobin A1c and body weight, but their long-term advantage over balanced dietary approaches was less consistent. High-fiber and whole-grain patterns were associated with lower T2D risk and improved postprandial metabolic responses. Protein source also appeared important, with plant-forward dietary patterns showing more favorable associations with insulin resistance than animal-heavy dietary patterns. Lifestyle-based programs and digital interventions improved adherence and self-management when supported by structured follow-up. The principal findings are summarized in Table 1.

Discussion

This systematic review supports the central role of nutrition in both prevention and management of T2D. The most consistent message across the included literature is that sustainable dietary patterns emphasizing whole foods, fiber-rich carbohydrates, unsaturated fats, and plant-forward choices are more clinically useful than rigid single-nutrient prescriptions. This interpretation is consistent with contemporary diabetes nutrition guidance, which emphasizes individualization, metabolic goals, patient preferences, comorbidities, and long-term adherence [8].

Mediterranean and plant-forward dietary patterns showed the most coherent benefits across glycemic and cardiometabolic outcomes. These patterns combine several favorable characteristics, including high intake of vegetables, legumes, whole grains, nuts, olive oil or other unsaturated fat sources, and relatively low intake of refined carbohydrates and ultra-processed foods. Their benefits are likely mediated through multiple mechanisms, including improved insulin sensitivity, reduced adiposity, lower systemic inflammation, and improved lipid profiles.

Low-carbohydrate dietary approaches may be useful for selected patients, particularly in the short term and when medication adjustment and clinical monitoring are available. However, long-term adherence, nutritional adequacy, lipid response, renal considerations, and patient acceptability should be considered before recommending highly restrictive approaches. The evidence does not support a single universal diet for all

patients with T2D. Instead, it supports individualized dietary planning that prioritizes safety, feasibility, cultural acceptability, and durability.

Fiber and whole grains represent another important component of diabetes prevention and management. Higher intake of dietary fiber and whole-grain foods may improve satiety, reduce postprandial glycemic excursions, enhance gut microbiota-related metabolic pathways, and support weight control. These findings argue against broadly discouraging all carbohydrate sources and instead support differentiating high-quality, minimally processed carbohydrates from refined starches and added sugars.

Protein source and overall dietary context also deserve attention. Evidence linking branched-chain amino acid profiles, insulin resistance, and animal-heavy dietary patterns suggests that protein quality may influence metabolic risk. Nevertheless, protein recommendations should be individualized according to kidney function, age, sarcopenia risk, weight goals, and comorbid disease burden.

This review has limitations. The included studies were heterogeneous in design, follow-up duration, dietary definitions, adherence assessment, and outcome reporting. Some dietary categories overlapped, and several interventions combined diet with physical activity, behavioral support, or digital tools, limiting attribution of effects to diet alone. The submitted manuscript also did not provide detailed study-level risk-of-bias ratings, which limits the ability to judge the certainty of evidence across categories.

Despite these limitations, the overall evidence is clinically practical. Nutritional care for T2D should not be framed as a short-term diet prescription but as a long-term therapeutic strategy integrated with weight management, medication review, behavioral counseling, and cardiometabolic risk reduction.

Conclusion

Nutrition plays a decisive role in T2D prevention and management. The current evidence favors individualized, sustainable, high-quality dietary patterns rather than a single rigid diet model. Mediterranean, plant-forward, fiber-rich, and minimally processed dietary patterns appear most consistently beneficial, while carbohydrate-restricted approaches may be useful in selected patients with appropriate clinical monitoring. Future research should focus on long-term adherence, implementation in routine care, cultural adaptation, and patient-centered outcomes.

References

- NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in diabetes prevalence and treatment from 1990 to 2022: a pooled analysis of 1108 population-representative studies with 141 million participants. *Lancet*. 2024;404(10467):2077-93. doi: 10.1016/S0140-6736(24)02317-1.
- GBD 2021 Diabetes Collaborators. Global, regional, and national burden of diabetes from 1990 to 2021, with projections of prevalence to 2050: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet*. 2023;402(10397):203-34. doi: 10.1016/S0140-6736(23)01301-6.
- Centers for Disease Control and Prevention. National Diabetes Statistics Report. Atlanta (GA): Centers for Disease Control and Prevention; 2024.
- World Health Organization. Diabetes. Geneva: World Health Organization; 2024. Available from: <https://www.who.int/health-topics/diabetes>
- Ajala O, English P, Pinkney J. Systematic review and meta-analysis of different dietary approaches to the management of type 2 diabetes. *Am J Clin Nutr*. 2013;97(3):505-16. doi: 10.3945/ajcn.112.042457.
- Viguiouk E, Stewart SE, Jayalath VH, Ng AP, Mirrahimi A, de Souza RJ, et al. Effect of vegetarian dietary patterns on cardiometabolic risk factors in diabetes: a systematic review and meta-analysis of randomized controlled trials. *Clin Nutr*. 2019;38(3):1133-45. doi: 10.1016/j.clnu.2018.05.032.
- Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med*. 2009;360(9):859-73. doi: 10.1056/NEJMoa0804748.
- Evert AB, Dennison M, Gardner CD, Garvey WT, Lau KHK, MacLeod J, et al. Nutrition therapy for adults with diabetes or prediabetes: a consensus report. *Diabetes Care*. 2019;42(5):731-54. doi: 10.2337/dci19-0014.
- Dinu M, Abbate R, Gensini GF, Casini A, Sofi F. Vegetarian, vegan diets and multiple health outcomes: a systematic review with meta-analysis of observational studies. *Crit Rev Food Sci Nutr*. 2017;57(17):3640-9. doi: 10.1080/10408398.2016.1138447.
- Tello-Montoliu A, Patel JV, Lip GYH. Mediterranean diet for the prevention of type 2 diabetes. *J Am Coll Cardiol*. 2019;73(24):3209-10.
- Bertoia ML, Mukamal KJ, Cahill LE, Hou T, Ludwig DS, Mozaffarian D, et al. Changes in intake of fruits and vegetables and weight change in United States men and women followed for up to 24 years: analysis from three prospective cohort studies. *PLoS Med*. 2015;12(9):e1001878. doi: 10.1371/journal.pmed.1001878.
- Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, et al. Primary prevention of cardiovascular disease with a Mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med*. 2018;378(25):e34. doi: 10.1056/NEJMoa1800389.
- Ros E, Singh A, O'Keefe JH. Nuts: natural pleiotropic nutraceuticals. *Nutrients*. 2021;13(9):3269. doi: 10.3390/nu13093269.
- Barnard ND, Cohen J, Jenkins DJA, Turner-McGrievy G, Gloede L, Green AA, et al. A low-fat vegan diet and a conventional diabetes diet in the treatment of type 2 diabetes: a randomized, controlled, 74-week clinical trial. *Am J Clin Nutr*. 2009;89(5):1588S-96S. doi: 10.3945/ajcn.2009.26736H.
- Ge L, Sadeghirad B, Ball GDC, da Costa BR, Hitchcock CL, Svendrovski A, et al. Comparison of dietary macronutrient patterns of 14 popular named dietary programmes for weight and cardiovascular risk factor reduction in adults: systematic review and network meta-analysis of randomised trials. *BMJ*. 2020;369:m696. doi: 10.1136/bmj.m696.
- Tay J, Thompson CH, Luscombe-Marsh ND, Wycherley TP, Noakes M, Buckley JD, et al. Effects of an energy-restricted low-carbohydrate, high-unsaturated fat/low-saturated fat diet versus a high-carbohydrate, low-fat diet in type 2 diabetes: a 2-year randomized clinical trial. *Am J Clin Nutr*. 2015;102(4):780-90. doi: 10.3945/ajcn.115.112581.
- Gardner CD, Trepanowski JF, Del Gobbo LC, Hauser ME, Rigdon J, Ioannidis JPA, et al. Effect of low-fat vs low-carbohydrate diet on 12-month weight loss in overweight adults and the association with genotype pattern or insulin secretion: the DIETFITS randomized clinical trial. *JAMA*. 2018;319(7):667-79. doi: 10.1001/jama.2018.0245.
- Westman EC, Yancy WS Jr, Humphreys M. Dietary treatment of diabetes mellitus in the pre-insulin era (1914-1922). *Diabetes Care*. 2019;42(7):1102-8. doi: 10.2337/dc18-2467.
- Goldenberg JZ, Day A, Brinkworth GD, Sato J, Yamada S, Jönsson T, et al. Efficacy and safety of low and very low carbohydrate diets for type 2 diabetes remission: systematic review and meta-analysis of published and unpublished randomized trial data. *BMJ*. 2021;372:m4743. doi: 10.1136/bmj.m4743.
- Meng Y, Bai H, Wang S, Li Z, Wang Q, Chen L. Efficacy of low carbohydrate diet for type 2 diabetes mellitus management: a systematic review and meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract*. 2017;131:124-31. doi: 10.1016/j.diabres.2017.07.006.
- Oh R, Gilani B, Uppaluri KR. Low-carbohydrate diet. *Am Fam Physician*. 2022;106(6):638-45.
- Harvard T.H. Chan School of Public Health. Carbohydrates and blood sugar. Boston (MA): Harvard T.H. Chan School of Public Health; 2024. Available from: <https://www.hsph.harvard.edu/nutritionsource/carbohydrates/carbohydrates-and-blood-sugar/>
- Chanson-Rolle A, Meynier A, Aubin F, Lappi J, Poutanen K, Vinoy S, et al. Systematic review and meta-analysis of human studies to support a quantitative recommendation for whole grain intake in relation to type 2 diabetes. *PLoS One*. 2015;10(6):e0131377. doi: 10.1371/journal.pone.0131377.
- InterAct Consortium. Dietary fibre and incidence of type 2 diabetes in eight European countries: the EPIC-InterAct Study and a meta-analysis of prospective studies. *Diabetologia*. 2015;58(7):1394-408. doi: 10.1007/s00125-015-3585-9.
- Aune D, Norat T, Romundstad P, Vatten LJ. Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Eur J Epidemiol*. 2013;28(11):845-58. doi: 10.1007/s10654-013-9852-5.
- Hu Y, Ding M, Sampson L, Willett WC, Manson JE, Wang M, et al. Intake of whole grain foods and risk of type 2 diabetes: results from three prospective cohort studies. *BMJ*. 2020;370:m2206. doi: 10.1136/bmj.m2206.
- McRae MP. Dietary fiber intake and type 2 diabetes mellitus: an umbrella review of meta-analyses. *J Chiropr Med*. 2018;17(1):44-53. doi: 10.1016/j.jcm.2017.11.002.
- Newgard CB, An J, Bain JR, Muehlbauer MJ, Stevens RD, Lien LF, et al. A branched-chain amino acid-related metabolic signature that differentiates obese and lean humans and contributes to insulin resistance. *Diabetes*. 2009;58(12):2791-9. doi: 10.2337/db09-0580.
- Chen Z, Zuurmond MG, van der Schaft N, Nano J, Wijnhoven HAH, Ikram MA, et al. Plant versus animal based diets and insulin resistance, prediabetes and type 2 diabetes: the Rotterdam Study. *Eur J Epidemiol*. 2018;33(9):883-93. doi: 10.1007/s10654-018-0414-8.
- Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002;346(6):393-403. doi: 10.1056/NEJMoa012512.
- Diabetes Prevention Program Research Group, Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet*. 2009;374(9702):1677-86. doi: 10.1016/S0140-6736(09)61457-4.
- U.S. Preventive Services Task Force, Curry SJ, Krist AH, Owens DK, Barry MJ, Caughey AB, et al. Behavioral counseling interventions to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;320(16):1677-87. doi: 10.1001/jama.2018.16789.
- Aziz Z, Absetz P, Oldroyd J, Pronk NP, Oldenburg B. A systematic review of real-world diabetes prevention programs: learnings from the last 15 years. *Implement Sci*. 2015;10:172. doi: 10.1186/s13012-015-0354-6.

Disclaimer/Publisher's Note: The statements, opinions, and data presented in publications in the Journal of Surgery and Medicine (JOSAM) are exclusively those of the individual author(s) and contributor(s) and do not necessarily reflect the views of JOSAM, the publisher, or the editor(s). JOSAM, the publisher, and the editor(s) disclaim any liability for any harm to individuals or damage to property that may arise from implementing any ideas, methods, instructions, or products referenced within the content. Authors are responsible for all content in their article(s), including the accuracy of facts, statements, and citations. Authors are responsible for obtaining permission from the previous publisher or copyright holder if re-using any part of a paper (e.g., figures) published elsewhere. The publisher, editors, and their respective employees are not responsible or liable for the use of any potentially inaccurate or misleading data, opinions, or information contained within the articles on the journal's website.